

ON THE LOCATION OF THE RITZ VALUES IN THE ARNOLDI PROCESS

GÉRARD MEURANT*

Abstract.

In this paper we give a necessary and sufficient condition for a set of complex values $\theta_1, \dots, \theta_k$ to be the Arnoldi Ritz values at iteration k for a general diagonalizable matrix A . Then we consider normal matrices and particularly real normal matrices with a real starting vector. We study in details the case $k = 2$ for which we characterize the boundary of the region in the complex plane where pairs of complex conjugate Ritz values are located. Several examples of computation of the boundary of the feasible region are given. Finally we formulate some conjectures and open problems for the location of the Arnoldi Ritz values in the case $k > 2$ for real normal matrices.

AMS subject classifications. 65F15, 65F18, 15A18

Key words. Arnoldi algorithm, eigenvalues, Ritz values, normal matrices.

1. Introduction. Approximations to (a few of) the eigenvalues (and eigenvectors) of large sparse non-Hermitian matrices are often computed with (variants of) the Arnoldi process. One of the most popular software packages is ARPACK [13]. It is used, for instance, in the Matlab function `eigs`. It uses the Implicitly Restarted Arnoldi algorithm. In this paper we are concerned with the standard Arnoldi process which for a matrix A of order n and a starting vector v (assumed to be of unit norm) computes a unitary matrix V with columns v^i and $v^1 = Ve_1 = v$ and an upper Hessenberg matrix H with positive real subdiagonal entries $h_{j+1,j}$, $j = 1, \dots, n-1$ such that

$$AV = VH,$$

if it does not stop before iteration n . The approximations of the eigenvalues of A (called the Ritz values) at step k are the eigenvalues $\theta_i^{(k)}$ of H_k , the leading principal submatrix of order k of H . The approximate eigenvectors are $x^i = V_{n,k}z_i^{(k)}$ where $z_i^{(k)}$ is the eigenvector associated with $\theta_i^{(k)}$ and $V_{n,k}$ is the matrix of the first k columns of V . In the sequel we will mainly consider the step k , so we will sometimes drop the upper index (k) . The relation satisfied by $V_{n,k}$ is

$$AV_{n,k} = V_{n,k}H_k + h_{k+1,k}v^{k+1}e_k^T,$$

where e_k is the last column of the identity matrix of order k . This equation shows how to compute the next column v^{k+1} of the matrix V and the k th column of H . When A is symmetric or Hermitian the Arnoldi process reduces to the Lanczos algorithm in which the matrix H is a symmetric tridiagonal matrix. There are many results about the convergence of the Lanczos Ritz values in the literature (see for instance [21], [17], [18], [19]). Most results are based on the Cauchy interlacing theorem which says that the Ritz values satisfy

$$\theta_1^{(k+1)} < \theta_1^{(k)} < \theta_2^{(k+1)} < \theta_2^{(k)} < \dots < \theta_k^{(k)} < \theta_{k+1}^{(k+1)},$$

*30 rue du sergent Bauchat, 75012 Paris, France (gerard.meurant@gmail.com). Started in Saint Aubin sur mer, August 2013, version September 27, 2013

and they are related to the eigenvalues λ_j by

$$\lambda_j < \theta_j^{(k)}, \quad \theta_{k+1-j}^{(k)} < \lambda_{n+1-j}, \quad 1 \leq j \leq k.$$

It is generally admitted that the convergence of the Lanczos process for Hermitian matrices is well understood. Unfortunately not much is known about the convergence of the Ritz values in the non-Hermitian case. Some results are known about the eigenvectors; see for instance [1], [2]. In fact the Arnoldi process may even not converge at all before the very last iteration. One can construct matrices with a given spectrum and starting vectors such that the Ritz values at all iterations are prescribed at arbitrary locations; see [9]. It means that we can construct examples for which the Ritz values *do not* converge to the eigenvalues of A before the last step.

However the matrices that can be build using this result have generally bad mathematical properties. In particular, they are not normal. In many practical cases we *do* observe convergence of the Ritz values toward eigenvalues. To try to understand the convergence when it occurs, an interesting problem is to know where are the locations of the Ritz values for a given matrix, in particular for matrices with special properties like (real) normal matrices. Of course, it is well known that they are inside the field of values of A which is defined as

$$W(A) = \{\theta \mid \theta = v^* A v, v \in \mathbb{C}^n, \|v\| = 1\}.$$

If the matrix A is normal the field of values is the convex hull of the eigenvalues and if the matrix is real it is symmetric with respect to the real axis.

The inverse problem described in Carden's thesis [4] and the paper [7] is, given a matrix A and complex values $\theta_1, \dots, \theta_k$, to know if there is a subspace of dimension k such that the θ_i 's are the corresponding Ritz values. If we restrict ourselves to Krylov subspaces and the Arnoldi algorithm, this amounts to know if there is a unit vector v such that the values θ_i are Ritz values for the Krylov subspace

$$\mathcal{K}_k(A, v) = \text{span}(v, Av, \dots, A^{k-1}v).$$

A closely related problem has been considered for normal matrices by Bujanović [3]. He was interested in knowing what is the location of the other Ritz values if one fixes some of the Ritz values in the field of values of A . He gave a necessary and sufficient condition that tells if a set of k complex values can be the Ritz values of a given normal matrix. Carden and Hansen [7] also gave a condition that is equivalent to Bujanović's. For normal matrices and $k = n - 1$, see [14] and for general matrices, see [6].

In this paper we first give a necessary and sufficient condition for a set of complex values $\theta_1, \dots, \theta_k$ to be the Arnoldi Ritz values at iteration k for a given general diagonalizable matrix A . This generalizes Bujanovic's condition. Then we restrict ourselves to real normal matrices and real starting vectors. We particularly study the case $k = 2$ for which we characterize the boundary of the region in the complex plane contained in $W(A)$ where pairs of complex conjugate Ritz values are located. We give several examples of computation of the boundary with real normal matrices of order up to 8. Finally, after describing some numerical experiments with real random starting vectors, we state some conjectures and open problems for $k > 2$ for real normal matrices. The aim of this section with only numerical results is to motivate other researchers for looking at these problems.

The paper is organized as follows. In section 2 we study the matrices H_k and we characterize the coefficients of their characteristic polynomial. Section 3 gives expressions for the entries of the matrix $M = K^*K$ where K is the Krylov matrix as a function of the eigenvalues and eigenvectors of A for diagonalizable matrices. This is used in section 4 where we give a necessary and sufficient condition for a set of k complex numbers to be the Arnoldi Ritz values at iteration k for diagonalizable matrices. The particular case of normal matrices is studied in section 5. The case of A real normal and $k = 2$ is considered in section 6 in which we characterize the boundary of the region where pairs of complex conjugate Ritz values are located. Open problems and conjectures for $k > 2$ and real normal matrices are described in section 7. Finally we give some conclusions.

2. The matrix H_k and the Ritz values. In this section, since the Ritz values are the eigenvalues of H_k , we are interested in characterizing the matrix H_k and the coefficients of its characteristic polynomial. In [15], [9], it was proved that the matrix H can be written as $H = UCU^{-1}$ where U is a nonsingular upper triangular matrix such that $K = VU$ with $K = (v \quad Av \quad \cdots \quad A^{n-1}v)$ and C is the companion matrix corresponding to the eigenvalues of A . We have the following theorem which characterizes H_k as a function of the entries of U .

THEOREM 2.1 ([10]). *For $k < n$ the matrix H_k can be written as $H_k = U_k C^{(k)} U_k^{-1}$, U_k being upper triangular and the principal submatrix of order k of U and $C^{(k)} = E_k + (0 \quad U_k^{-1} U_{[1:k],k+1})$, a companion matrix where E_k is a square down-shift matrix of order k ,*

$$E_k = \begin{pmatrix} 0 & & & & \\ 1 & 0 & & & \\ & \ddots & \ddots & & \\ & & & 1 & 0 \\ & & & & 1 & 0 \end{pmatrix}.$$

Moreover the subdiagonal entries of H are $h_{j+1,j} = \frac{U_{j+1,j+1}}{U_{j,j}}$, $j = 1, \dots, n-1$.

Clearly the Ritz values at step k are the eigenvalues of $C^{(k)}$. We see that they only depend on the matrix U and its inverse. They are also the roots of the monic polynomial defined below. By considering at what is the inverse of an upper triangular matrix we note that the last column of $C^{(k)}$ can be written as

$$U_k^{-1} U_{[1:k],k+1} = -U_{k+1,k+1} (U_{k+1}^{-1})_{[1:k],k+1} = -U_{k+1,k+1} (U^{-1})_{[1:k],k+1}.$$

Hence, up to a multiplying coefficient, the last column of $C^{(k)}$ is made of the k first components of the $k+1$ st column of the inverse of U . The last column of $C^{(k)}$ gives the coefficients of the characteristic polynomial of H_k . Let

$$\begin{pmatrix} \beta_0^{(k)} \\ \vdots \\ \beta_{k-1}^{(k)} \end{pmatrix} = -U_k^{-1} U_{[1:k],k+1}.$$

The Ritz values are the roots of the polynomial $q_k(\lambda) = \lambda^k + \sum_{j=0}^{k-1} \beta_j^{(k)} \lambda^j = \prod_{i=1}^k (\lambda - \theta_i^{(k)})$. Since the entries of U and U^{-1} are intricate functions of the eigenvalues and

eigenvectors of A , the following theorem provides a simpler characterization of the coefficients of the characteristic polynomial of H_k .

THEOREM 2.2. *The vector of the coefficients of the characteristic polynomial of H_k denoted as $(\beta_0^{(k)} \ \dots \ \beta_{k-1}^{(k)})$ is the solution of the linear system,*

$$(2.1) \quad M_k \begin{pmatrix} \beta_0^{(k)} \\ \vdots \\ \beta_{k-1}^{(k)} \end{pmatrix} = -M_{[1:k],k+1},$$

where $M_k = U_k^* U_k$.

Proof. From what we have seen above the proof is straightforward. We have

$$U_k \begin{pmatrix} \beta_0^{(k)} \\ \vdots \\ \beta_{k-1}^{(k)} \end{pmatrix} = -U_{[1:k],k+1}.$$

Multiplying by U_k^* we obtain

$$M_k \begin{pmatrix} \beta_0^{(k)} \\ \vdots \\ \beta_{k-1}^{(k)} \end{pmatrix} = -U_k^* U_{[1:k],k+1}.$$

Clearly $U_k^* U_{[1:k],k+1} = M_{[1:k],k+1}$. \square

Therefore it is interesting to consider the matrix $M = U^* U$ and its principal submatrices. This is done in the next section.

3. The matrix M . In this section we characterize the entries of $M = U^* U = K^* K$ as functions of the eigenvalues and eigenvectors of A and of the starting vector v for diagonalizable matrices A .

THEOREM 3.1. *Let the spectral decomposition of A be $A = X \Lambda X^{-1}$ with eigenvalues λ_i , $i = 1, \dots, n$. The entries of $M = U^* U$ are given by*

$$(3.1) \quad M_{\ell,m} = \sum_{i=1}^n \sum_{j=1}^n (X^* X)_{i,j} \bar{c}_i c_j \bar{\lambda}_i^{\ell-1} \lambda_j^{m-1}, \quad \ell, m = 1, \dots, n,$$

with $c = X^{-1}v$. If the matrix A is normal we have a simpler expression,

$$(3.2) \quad M_{\ell,m} = \sum_{i=1}^n |c_i|^2 \bar{\lambda}_i^{\ell-1} \lambda_i^{m-1}, \quad \ell, m = 1, \dots, n,$$

with $c = X^* v$.

Proof. Since we have assumed that the matrix A is diagonalizable with eigenvalues λ_i we have

$$K = X (c \ \Lambda c \ \dots \ \Lambda^{n-1} c),$$

where $c = X^{-1}v$. Let D_c be the diagonal matrix with diagonal entries c_j . The matrix K is

$$K = X D_c \mathcal{V},$$

with the Vandermonde matrix,

$$\mathcal{V} = \begin{pmatrix} 1 & \lambda_1 & \cdots & \lambda_1^{n-1} \\ 1 & \lambda_2 & \cdots & \lambda_2^{n-1} \\ \vdots & \vdots & & \vdots \\ 1 & \lambda_n & \cdots & \lambda_n^{n-1} \end{pmatrix}.$$

We note that this factorization of the Krylov matrix has been used in [11]; see also [22]. Therefore $M = K^*K = \mathcal{V}^*D_{\bar{c}}X^*XD_c\mathcal{V}$. If A is normal, $X^*X = I$ and $M = \mathcal{V}^*D_{\omega}\mathcal{V}$ with $\omega_j = |c_j|^2$. The entries of M can be obtained as a function of the eigenvalues and eigenvectors of A by

$$\begin{aligned} M_{\ell,m} &= e_{\ell}^T M e_m \\ &= e_{\ell}^T \mathcal{V}^* D_{\bar{c}} X^* X D_c \mathcal{V} e_m \\ &= (\bar{\lambda}_1^{\ell-1} \quad \cdots \quad \bar{\lambda}_n^{\ell-1}) D_{\bar{c}} X^* X D_c \begin{pmatrix} \lambda_1^{m-1} \\ \vdots \\ \lambda_n^{m-1} \end{pmatrix} \\ &= \sum_{i=1}^n \sum_{j=1}^n (X^* X)_{i,j} \bar{c}_i c_j \bar{\lambda}_i^{\ell-1} \lambda_j^{m-1}. \end{aligned}$$

If A is normal, we have $X^*X = I$ and

$$M_{\ell,m} = \sum_{i=1}^n |c_i|^2 \bar{\lambda}_i^{\ell-1} \lambda_i^{m-1}.$$

This last result was already known from [20]. \square

4. The inverse problem for diagonalizable matrices. For the first Arnoldi iteration (that is, $k = 1$) the inverse problem is always solvable. We have $h_{1,1} = v^*Av$. For $\theta^{(1)} \in W(A)$ there exists a vector v such that $\theta^{(1)} = v^*Av$. Algorithms for computing such vectors are given in [5], [8], [16]. We note that if A and v are real, the first Ritz value $\theta_1^{(1)}$ is real.

For the inverse problem at Arnoldi iteration $k > 1$ we assume that we have a set of k given complex numbers $\theta_1, \dots, \theta_k$ belonging to $W(A)$ and we would like to find (if possible) a vector v of unit norm such that the θ_j 's are the Ritz values at iteration k when running the Arnoldi algorithm with (A, v) .

From (2.1) we have an equation relating the coefficients of the characteristic polynomial of H_k and the entries of a submatrix of the matrix M . Since the Ritz values are zeros of the polynomial $\lambda^k + \sum_{j=0}^{k-1} \beta_j^{(k)} \lambda^j = \prod_{i=1}^k (\lambda - \theta_i)$, the coefficients $\beta_j^{(k)}$ are (up to the sign) elementary symmetric functions of the θ_j 's. Therefore

$$(4.1) \quad \beta_j^{(k)} = (-1)^{k-j} e_{k-j}(\theta_1, \dots, \theta_k), \quad j = 0, \dots, k-1,$$

with

$$e_i(\theta_1, \dots, \theta_k) = \sum_{1 \leq j_1 < j_2 < \cdots < j_i \leq k} \theta_{j_1} \cdots \theta_{j_i}, \quad i = 1, \dots, k.$$

Therefore we have the following characterization of the existence of a starting vector.

THEOREM 4.1. *There exists a starting vector $v = Xc$ of unit norm such that $\theta_1, \dots, \theta_k$ are the Arnoldi Ritz values at iteration k if and only if the nonlinear system (2.1) with unknowns c_j , $j = 1, \dots, n$ (where the coefficients $\beta_j^{(k)}$ are defined by (4.1)) to which we add*

$$(4.2) \quad \sum_{i,j=1}^n \bar{c}_i c_j (X^* X)_{i,j} = 1,$$

has at least one solution vector c .

Proof. Let us assume that there exists a vector v such that $\theta_1, \dots, \theta_k$ are the Arnoldi Ritz values at iteration k . They are the roots of the characteristic polynomial whose coefficients $\beta_j^{(k)}$ are given by (4.1). Hence by Theorem 2.2 the coefficients are solution of the linear system (2.1) and the vector c is a solution of the nonlinear system defined by (2.1) plus (4.2) because the vector v is of unit norm.

Conversely if there is a solution c to the nonlinear system (2.1)-(4.2) then there exists a solution of the linear system (2.1) with unknowns $\beta_j^{(k)}$ which by Theorem 2.2 are the coefficients of the characteristic polynomial of H_k and the complex numbers defined as the roots of the polynomial are the Ritz values at Arnoldi iteration k . \square

To make the things clear let us consider the case $k = 2$ with $\theta_1 = \theta_1^{(2)}$, $\theta_2 = \theta_2^{(2)}$ given. Let $p = \theta_1 \theta_2$ and $s = \theta_1 + \theta_2$ which are known. We note that M_2 is an Hermitian matrix. Then (2.1) is

$$M_2 \begin{pmatrix} p \\ -s \end{pmatrix} = -M_{[1:2],3}.$$

Therefore we have the two equations,

$$p - sM_{1,2} = -M_{1,3}, \quad sM_{2,2} = M_{2,3} + p\overline{M_{1,2}}.$$

The equations to be satisfied are

$$p - s \sum_{i,j=1}^n \bar{c}_i c_j (X^* X)_{i,j} \lambda_j = - \sum_{i,j=1}^n \bar{c}_i c_j (X^* X)_{i,j} \lambda_j^2,$$

$$s \sum_{i,j=1}^n \bar{c}_i c_j (X^* X)_{i,j} \bar{\lambda}_i \lambda_j = \sum_{i,j=1}^n (X^* X)_{i,j} \bar{c}_i c_j \bar{\lambda}_i \lambda_j^2 + p \sum_{i,j=1}^n (X^* X)_{i,j} \bar{c}_i c_j \bar{\lambda}_i.$$

Since we need to find a vector v of unit norm, we have to add the equation $\|Xc\|^2 = c^* X^* X c = 1$. It yields

$$\sum_{i,j=1}^n \bar{c}_i c_j (X^* X)_{i,j} = 1.$$

Because we know s and p , this is three nonlinear complex equations in n complex unknowns c_i , $i = 1, \dots, n$. Whether or not this system has solutions determines if θ_1 and θ_2 are feasible values since, if a solution c exists, we can then find a vector v such that the two given values θ_1 and θ_2 are Ritz values for $\mathcal{K}_2(A, v)$.

We remark that this is generally not a polynomial system because of the conjugacy in $\bar{c}_i c_j$. However, we can convert this system to a polynomial system by considering

the real and imaginary parts of c_i as unknowns. We have then a polynomial system of six equations in $2n$ unknowns with complex coefficients that can be converted to a polynomial system with real coefficients by taking the real and imaginary parts. The trouble then is that we have to know if there are real solutions. Unfortunately there are not many results about this problem in the algebraic geometry literature. The situation is much simpler if we assume that the matrix A is normal. This case is considered in the next section.

5. The inverse problem for normal matrices. For a normal matrix and assuming that we know the coefficients $\beta_0^{(k)}, \dots, \beta_{k-1}^{(k)}$, we obtain a $(k+1) \times n$ linear system for the moduli squared $\omega_i = |c_i|^2$. It yields a linear system

$$(5.1) \quad C_C \omega = f_C.$$

If we put the normalizing equation $\sum_{i=1}^n \omega_i = 1$ first, the entries of C_C are all 1 for the first row. The entries of the second row are

$$(C_C)_{2,m} = \sum_{i=1}^{k-1} \beta_i^{(k)} \lambda_m^i + \lambda_m^k, \quad m = 1, \dots, n,$$

and the other entries are

$$(C_C)_{\ell,m} = \sum_{i=0}^{k-1} \beta_i^{(k)} \bar{\lambda}_m^{\ell-2} \lambda_m^i + \bar{\lambda}_m^{\ell-2} \lambda_m^k, \quad \ell = 3, \dots, k+1, \quad m = 1, \dots, n.$$

The right-hand side is all zero, except for the first two components, $(f_C)_1 = 1$, $(f_C)_2 = -\beta_0^{(k)}$.

We can also turn this linear system of $k+1$ complex equations in n real unknowns into a real linear system by taking the real and imaginary parts of rows 2 to k . It gives a $(2k+1) \times n$ matrix C_R and the right-hand side is zero except for the three first components $(f_R)_1 = 1$, $(f_R)_2 = -\text{Re}[\beta_0^{(k)}]$, $(f_R)_3 = -\text{Im}[\beta_0^{(k)}]$.

Compared to the case of a general diagonalizable matrix studied in the previous section there are good things and bad things. The good thing is that we have a linear system for the unknowns ω_i instead of a nonlinear one. The bad thing is that we need to find a solution which is real and positive. Having (if possible) a real solution is easy by solving $C_R \omega = f_R$ but we still need a positive solution. The characterization of $\theta_1, \dots, \theta_k$ being feasible is given in the following theorem.

THEOREM 5.1. *Let A be a normal matrix. There exists a starting vector $v = Xc$ of unit norm such that $\theta_1, \dots, \theta_k$ are the Arnoldi Ritz values at iteration k if and only if the linear system $C_R \omega = f_R$ where the coefficients $\beta_j^{(k)}$ are defined by (4.1) has at least one solution vector ω with $\omega_i \geq 0$, $i = 1, \dots, n$. Then c is any vector such that $|c_i|^2 = \omega_i$.*

Proof. The proof is similar to the proof of Theorem 4.1. \square

The condition given in Theorem 5.1 must be equivalent to the condition recently proposed by Bujanović (Theorem 4 in [3]).

Things can be further simplified if the matrix A and the starting vector are real. Up to our knowledge this case has not been considered by other authors. Then the eigenvalues of A are real or occur in complex conjugate pairs. If the starting vector v is real, everything is real in the Arnoldi algorithm (in particular the matrix H) and the

Ritz values are real or complex conjugate pairs which are the roots of a polynomial with real coefficients $\beta_j^{(k)}$. The two eigenvectors of A corresponding to a complex conjugate pair are conjugate and the eigenvectors corresponding to real eigenvalues are real. Then, with v real, if $c = X^*v$ and $\lambda_i = \bar{\lambda}_j$ we have $c_i = \bar{c}_j$. This means that when the Ritz values are known, we have only one unknown c_i for each pair of complex conjugate eigenvalues. Let us assume that the matrix A has p_C pairs of complex conjugate eigenvalues (with $2p_C \leq n$) that are listed first and $n - 2p_C$ real eigenvalues denoted $(\lambda_{2p_C+1}, \dots, \lambda_n)$. Then, we have only $n - p_C$ unknowns that, to avoid some confusion, we denote with their initial indices ranging from 1 to n as for the eigenvalues. That is, the unknowns will be the components of the vector

$$(5.2) \quad \tilde{\omega} = (|c_1|^2, |c_3|^2, \dots, |c_{2p_C-1}|^2, |c_{2p_C+1}|^2, |c_{2p_C+2}|^2, \dots, |c_n|^2)^T.$$

Then, in the equations derived from the matrix M we have to group the terms containing λ_i and $\bar{\lambda}_i$. Since everything is real we denote the matrix as C_R even though it is different from the matrix described above. The first row of the matrix C_R is now

$$(5.3) \quad (C_R)_{1,m} = 2, \quad m = 1, \dots, p_C, \quad (C_R)_{1,m} = 1, \quad m = p_C + 1, \dots, n - p_C.$$

The second row is

$$(5.4) \quad (C_R)_{2,m} = 2 \sum_{i=1}^{k-1} \beta_i^{(k)} \operatorname{Re}(\lambda_{2m-1}^i) + 2\operatorname{Re}(\lambda_{2m-1}^k), \quad m = 1, \dots, p_C,$$

$$(5.5) \quad (C_R)_{2,m} = \sum_{i=1}^{k-1} \beta_i^{(k)} \lambda_{p_C+m}^i + \lambda_{p_C+m}^k, \quad m = p_C + 1, \dots, n - p_C,$$

and the other entries are

$$(5.6) \quad (C_R)_{\ell,m} = 2 \sum_{i=0}^{k-1} \beta_i^{(k)} \operatorname{Re}(\bar{\lambda}_{2m-1}^{\ell-2} \lambda_{2m-1}^i) + 2\operatorname{Re}(\bar{\lambda}_{2m-1}^{\ell-2} \lambda_{2m-1}^k), \quad \ell = 3, \dots, k+1, \quad m = 1, \dots, p_C,$$

$$(5.7) \quad (C_R)_{\ell,m} = \sum_{i=0}^{k-1} \beta_i^{(k)} \lambda_{p_C+m}^{\ell-2+i} + \lambda_{p_C+m}^{\ell-2+k}, \quad \ell = 3, \dots, k+1, \quad m = p_C + 1, \dots, n - p_C.$$

The right-hand side is all zero, except for the first two components, $(f_R)_1 = 1$, $(f_R)_2 = -\beta_0^{(k)}$. Therefore, the real matrix C_R is only $(k+1) \times (n - p_C)$ and we have $n - p_C$ unknowns. To know if there exist a positive solution we have to consider the cases $k+1 > n - p_C$ (overdetermined system), $k+1 = n - p_C$ (square system) and $k+1 < n - p_C$ (underdetermined system). When we have a positive solution we can find a real vector c by expanding the solution and taking square roots and finally a real starting vector $v = Xc$. The previous discussion is summarized in the following theorem.

THEOREM 5.2. *Let A be a real normal matrix. There exists a real starting vector $v = Xc$ of unit norm such that $\theta_1, \dots, \theta_k$, where these values are real or occur in complex conjugate pairs, are the Arnoldi Ritz values at iteration k if and only if the linear system $C_R \tilde{\omega} = f_R$, where the coefficients $\beta_j^{(k)}$ are defined by (4.1), the matrix*

C_R is defined by (5.3)-(5.7) and $\tilde{\omega}$ by (5.2), has at least one solution vector $\tilde{\omega}$ with $\tilde{\omega}_i \geq 0$, $i = 1, \dots, n - p_C$. Then c is any real vector such that $|c_i|^2 = \omega_i$ where ω is given by the expansion of $\tilde{\omega}$.

Let us now consider the problem of finding a positive solution in the case the linear system $C_R \tilde{\omega} = f_R$ is underdetermined that is, $k + 1 < n - p_C$. Solutions of a system like this can be found by using the Singular Value Decomposition (SVD). Let us consider the generic case where C_R has full rank $k + 1$. The matrix can be factorized as

$$C_R = \hat{U} \begin{pmatrix} D & 0 \end{pmatrix} \hat{V}^*, \quad D \text{ diagonal}, \quad \hat{U}^* \hat{U} = I, \quad \hat{V}^* \hat{V} = I.$$

The orthonormal matrix \hat{U} is of order $k + 1$ as well as D and \hat{V} is of order $n - p_C$. The diagonal of D contains the singular values. Since all the singular values are non-zero we can find solutions to

$$\hat{U} \begin{pmatrix} D & 0 \end{pmatrix} \hat{V}^* \tilde{\omega} = \begin{pmatrix} 1 \\ (f_R)_2 \\ 0 \end{pmatrix}.$$

Let $y = \hat{V}^* \tilde{\omega}$,

$$\hat{U} \begin{pmatrix} D & 0 \end{pmatrix} y = \begin{pmatrix} 1 \\ (f_R)_2 \\ 0 \end{pmatrix} \Rightarrow \hat{y} \equiv \begin{pmatrix} y_1 \\ \vdots \\ y_{k+1} \end{pmatrix} = D^{-1} \hat{U}^* \begin{pmatrix} 1 \\ (f_R)_2 \\ 0 \end{pmatrix}.$$

The solutions are given by

$$\tilde{\omega} = \hat{V} \begin{pmatrix} y_1 \\ \vdots \\ y_{k+1} \\ \times \\ \vdots \\ \times \end{pmatrix},$$

where the \times denote arbitrary real numbers. Let $\hat{V} = [\hat{V}_1 \ \hat{V}_2]$ with \hat{V}_1 having $k + 1$ columns. Then, we have a positive solution if and only if there exists a vector z such that

$$(5.8) \quad -\hat{V}_2 z \leq \hat{V}_1 \hat{y}$$

and

$$\tilde{\omega} = \hat{V} \begin{pmatrix} \hat{y} \\ z \end{pmatrix}.$$

To check if there is a solution to the system of inequalities (5.8) we use the algorithm described in [12] that was intended to convert a system of linear inequalities to a representation using the vertices of the polyhedron defined by the inequalities. It relies on computing the rank of submatrices and tell us if the system is feasible or not.

6. A real normal, $k = 2$. In this section we further simplify the problem and concentrate on the case $k = 2$ for a real normal matrix and a real starting vector. The matrix H_2 is real and has either two real eigenvalues or a pair of complex conjugate eigenvalues. We are interested in the latter case for which we have $\theta_2 = \bar{\theta}_1$. Hence it is enough to look for the location of the complex Ritz value θ_1 and this considerably simplifies the problem. We call the set of all the complex values θ_1 in the field of values yielding a positive solution *the feasible region*. To obtain a graphical representation of the feasible region we can proceed as in Bujanović [3]. We set up a regular cartesian mesh over the field of values (in fact over the smallest rectangle containing the upper part - $y \geq 0$ - of the field of values) of A for the values of θ_1 and we check if there are positive solutions to the $3 \times (n - p_C)$ linear system $C_R \tilde{\omega} = f_R$ for each value of $\theta_1 = (x, y)$ in the mesh by considering the system of inequalities (5.8). When the system is feasible for a given value of θ_1 on the mesh we flag this location. Hence, for each θ_1 in the marked area we can find a real vector v such that $\theta_1, \theta_2 = \bar{\theta}_1$ are the Ritz values at iteration 2. This gives an approximation of the feasible region. For θ_1 outside of the feasible region there does not exist a real vector v that yields $(\theta_1, \bar{\theta}_1)$ as Arnoldi Ritz values at iteration 2. Of course this way of looking for the feasible locations of θ_1 by discretizing the field of values has some drawbacks since some tiny feasible parts may be missing if the discretization is not fine enough.

Figures 6.1 and 6.2 display an example (Example 1) of a matrix of order 4 with two real eigenvalues on each side of the real part of λ_1 . In this example the matrix C_R is square of order 3 since $n - p_C = 4 - 1 = 3$ and nonsingular. The field of values is shown in red and the eigenvalues of A are the red circles. The feasible values of θ_1 (resp. θ_2) are marked with blue (resp. red) crosses. In this example the feasible region is a surface in the complex plane. In this case it is connected and convex (if we add the real values inside the region) but we will see later that this is not always the case. Of course, we can also have two real Ritz values outside this region. In this example it seems that the real Ritz values can be anywhere in the interval defined by the two real eigenvalues of A . Figure 6.2 was obtained by using 700 random real starting vectors and running the Arnoldi algorithm. We see that we obtain approximately the same shape for the feasible region. In Example 1 the matrix A is

$$A = \begin{pmatrix} -0.446075 & 0.358311 & -0.605655 & 1.12896 \\ -0.512738 & -0.263009 & -1.09795 & 0.285 \\ 1.15846 & 0.636041 & -0.72035 & 0.0184702 \\ -0.405993 & -1.00831 & -0.417846 & -0.456834 \end{pmatrix}.$$

The eigenvalues of A are

$$-0.432565 + 1.66558i, \quad -0.432565 - 1.66558i, \quad 0.187377, \quad -1.20852.$$

Since we may miss some parts of the feasible region due to a too coarse discretization, it is interesting to characterize its boundary. This can be done by explicitly writing down the inverse of the matrix C_R and looking at the inequalities given by the positivity constraints for ω_j . It corresponds to the elimination of the components of ω in the equations. For simplicity let us denote

$$C_R = \begin{pmatrix} 2 & 1 & 1 \\ a & c & e \\ b & d & f \end{pmatrix}.$$




FIG. 6.1. *Example 1, $n = 4, k = 2$, A normal real, location of $\theta_1 = \bar{\theta}_2$*




FIG. 6.2. *Example 1, $n = 4, k = 2$, A normal real, location of $\theta_1 = \bar{\theta}_2$, Arnoldi with random real vectors v*

The inverse is given by

$$C_R^{-1} = \frac{1}{D} \begin{pmatrix} cf - ed & d - f & e - c \\ eb - af & 2f - b & a - 2e \\ ad - cb & b - 2d & 2c - a \end{pmatrix}, \quad D = a(d - f) + c(2f - b) + e(b - 2d).$$

We apply the inverse to the right-hand side (which, after a change of signs, is $(1 \quad p \quad 0)^T$, $p = |\theta_1|^2$) and we get

$$\omega = \frac{1}{D} \begin{pmatrix} cf - ed + (d - f)p \\ eb - af + (2f - b)p \\ ad - cb + (b - 2d)p \end{pmatrix}.$$

We are interested in the components of ω being positive. The outside of the feasible region is characterized by the fact that at least one component ω_j is negative. Therefore the boundary must be given by some of the components of the solution being

zero. Hence, we have to look at the three equations

$$\begin{aligned} cf - ed + (d - f)p &= 0, \\ eb - af + (2f - b)p &= 0, \\ ad - cb + (b - 2d)p &= 0. \end{aligned}$$

The coefficients a, b, c, d, e, f are functions of unknowns quantities $s = 2x = 2\operatorname{Re}(\theta_1)$ and $p = x^2 + y^2 = |\theta_1|^2$. These equations define three curves in the (x, y) complex plane. Some (parts) of these curves yield the boundary of the feasible region for θ_1 . However we note that one component can be zero on one of the curves without changing sign. Therefore all the curves may not be relevant for the boundary. We just know that the boundary is contained in the union of the curves. Moreover we are only interested in the parts of the curves contained in the convex hull of the eigenvalues of A . For completeness, remember that we have

$$a = 2s\operatorname{Re}(\lambda_1) - 2\operatorname{Re}(\lambda_1^2), \quad b = 2s|\lambda_1|^2 - 2|\lambda_1|^2\operatorname{Re}(\lambda_1) - 2p\operatorname{Re}(\lambda_1),$$

$$c = s\lambda_3 - \lambda_3^2, \quad d = s\lambda_3^2 - \lambda_3^3 - p\lambda_3,$$

$$e = s\lambda_4 - \lambda_4^2, \quad f = s\lambda_4^2 - \lambda_4^3 - p\lambda_4.$$

The first curve involves only the real eigenvalues of A . The two other curves pass through λ_1 and $\bar{\lambda}_1$.

Figure 6.3 shows the boundary curves we obtain for Example 1 as well as the approximation of the feasible region using a smaller number of discretization points than before. These curves were obtained using a contour plot of level 0 for the three functions of x and y . We see that we do get the boundary of the feasible region for θ_1 . We have only two curves since the one which depends only on the real eigenvalues (corresponding to the first equation) does not have points in the window we are interested in (that is, the field of values).




FIG. 6.3. Example 1, $n = 4, k = 2$, A normal real, location of $\theta_1 = \bar{\theta}_2$ and the boundary of the feasible region

Let us now consider $n = 5$. The matrix A has either two pairs of complex conjugate eigenvalues $(\lambda_1, \bar{\lambda}_1)$, $(\lambda_3, \bar{\lambda}_3)$ and a real eigenvalue λ_4 (that can be on the boundary or inside the field of values) or one pair of complex conjugate eigenvalues and three real eigenvalues (there is at least one inside the field of values, eventually two). In the first case the matrix C_R is 3×3 as it was for the previous example with $n = 4$. Then, we can apply the same techniques by eliminating the components of ω to obtain the boundary of the feasible region for θ_1 . The only differences are that $(C_R)_{1,2} = 2$ and the values of the coefficients. We have

$$a = 2s\operatorname{Re}(\lambda_1) - 2\operatorname{Re}(\lambda_1^2), \quad b = 2s|\lambda_1|^2 - 2|\lambda_1|^2\operatorname{Re}(\lambda_1) - 2p\operatorname{Re}(\lambda_1),$$

$$c = 2s\operatorname{Re}(\lambda_3) - 2\operatorname{Re}(\lambda_3^2), \quad d = 2s|\lambda_3|^2 - 2|\lambda_3|^2\operatorname{Re}(\lambda_3) - 2p\operatorname{Re}(\lambda_3),$$

$$e = s\lambda_4 - \lambda_4^2, \quad f = s\lambda_4^2 - \lambda_4^3 - p\lambda_4.$$

The equations are

$$cf - ed + (d - 2f)p = 0,$$

$$eb - af + (2f - b)p = 0,$$

$$ad - bc + 2(b - d)p = 0.$$

This is only small changes from above with a multiplying coefficient 2 for some terms. Figure 6.4 displays the feasible region and the boundary for a case with two pairs of complex conjugate eigenvalues and a real eigenvalue inside the field of values (Example 2). In this example the feasible region is not connected nor convex. Figure 6.5 shows that random starting vectors do not always give a good rendering of the feasible region. In Example 2 the matrix is

$$A = \begin{pmatrix} 0.513786 & -0.419578 & 0.156842 & 0.447046 & 0.540983 \\ -0.789795 & 0.767537 & -0.451475 & 0.12333 & 0.202036 \\ 0.0825256 & -0.091751 & 1.31755 & 0.5561 & -0.00409194 \\ 0.179105 & 0.7687 & -0.247999 & 1.31189 & 0.0474895 \\ -0.174622 & -0.329046 & -0.185905 & 0.403025 & -0.101738 \end{pmatrix}.$$

The eigenvalues of A are

$$\begin{aligned} & -0.178102 + 0.498599i, & -0.178102 - 0.498599i, \\ & 1.5788 + 0.584391i, & 1.5788 - 0.584391i, & 1.00762. \end{aligned}$$

Figure 6.6 displays an example with only one pair of complex conjugate eigenvalues and three real eigenvalues and two of them on the same side of the real part of the complex eigenvalues (Example 3). We see that we have one piece of curve which is inside the feasible region. It can be considered as a “spurious” curve (even though we will see later on that these curves can have also some interest). The matrix of Example 3 is

$$A = \begin{pmatrix} -1.07214 & -0.549535 & 0.809383 & -0.0826907 & 0.345094 \\ -0.779134 & 1.06039 & 0.100179 & 0.621762 & -0.184854 \\ -0.33126 & -0.0693308 & -0.551724 & 1.39559 & 1.19566 \\ -0.24838 & 0.134568 & -0.902458 & -0.0781342 & -1.22051 \\ -0.551853 & -0.844358 & -1.41854 & 0.206828 & -0.233364 \end{pmatrix}.$$




FIG. 6.4. *Example 2, $n = 5, k = 2$, A normal real, location of $\theta_1 = \bar{\theta}_2$*




FIG. 6.5. *Example 2, $n = 5, k = 2$, A normal real, location of $\theta_1 = \bar{\theta}_2$, Arnoldi with random real vectors v*




FIG. 6.6. *Example 3, $n = 5, k = 2$, A normal real, location of $\theta_1 = \bar{\theta}_2$*

The eigenvalues of A are

$$-0.600433 + 2.06392i, \quad -0.600433 - 2.06392i, \quad -1.40594, \quad 1.56985, \quad 0.161981.$$

For n larger than 5 the situation is more complicated to compute the boundary. We generally have more than 3 unknowns (except for $n = 6$ with three pairs of complex conjugate eigenvalues) and therefore an underdetermined linear system for the ω_j 's. When prescribing a value of θ_1 (with $\theta_2 = \bar{\theta}_1$), as we have seen before we can check the feasibility by using the SVD of the rectangular matrix $C_R = \hat{U}S\hat{V}^T$.

Concerning the boundary of the feasible region, the pieces of the boundary correspond to some of the components of ω being zero. Therefore, we can apply the same elimination technique as before by considering the matrices of order 3 corresponding to all the triples of eigenvalues, a pair of complex conjugate ones counting only for one. It corresponds to considering only three components of ω putting the other components to zero. We have to consider 3×3 matrices similar as the ones we had before with a first row being $(2, 2, 2)$, $(2, 1, 1)$ or $(1, 1, 1)$. The number of curves is three times the number of triples of eigenvalues.

Doing this corresponds to what is done to handle the linear constraints in linear programming (LP) whose solution components must be positive. Let us assume that we have linear equality constraints $Cx = b$ defined by a real $m \times n$ matrix C of full rank with $m < n$. Possibly doing a permutation of columns we can write $C = [B \ E]$ with B nonsingular of order m . Then

$$x = \begin{pmatrix} B^{-1}b \\ 0 \end{pmatrix}$$

is called a basic solution. It is degenerate if some components of $B^{-1}b$ are zero. This is just taking m independent columns of C , putting the other components of the solution to zero and solving. A basic feasible solution (BFS) is a basic solution that satisfies the constraints of the LP. The feasible region defined by the constraints is convex, closed and bounded from below. The feasible region is a polyhedron and it can be shown that the BFS are extreme points (vertices or corners) of the feasible region.

This is similar to what we are doing. We have a polyhedron in the ω -space defined by the system with matrix C_R and considering all the 3×3 matrices (provided they are nonsingular) is computing symbolically the basic solutions. The feasible ones (with $\omega_j \geq 0$) correspond to some vertices of the polyhedron. Clearly these curves are where components of ω may change signs as a function of $x = \text{Re}(\theta_1)$ and $y = \text{Im}(\theta_1)$. They also give a parametric description of the vertices of the polyhedron.

Figure 6.7 corresponds to an example with $n = 6$ and three pairs of complex conjugate eigenvalues (Example 4). In this example the matrix C_R is square of order 3, ω has only three components and there is no spurious curve. We see that the shape of the feasible region can be quite complicated. The matrix A of Example 4 is

$$A = \begin{pmatrix} -0.401151 & 0.0951597 & 0.336817 & -0.0155421 & 0.342989 & 0.059462 \\ 0.0435544 & -0.711607 & -0.0851345 & -0.100931 & 0.19691 & -0.0848016 \\ -0.3473 & 0.0330741 & -0.458265 & 0.338473 & 0.161655 & -0.163792 \\ 0.0903354 & 0.144387 & 0.0427703 & -0.167152 & 0.14634 & -0.661259 \\ -0.309586 & -0.118264 & 0.148041 & -0.196687 & -0.517635 & -0.205145 \\ 0.131915 & -0.142526 & 0.380522 & 0.570822 & -0.0924846 & -0.165907 \end{pmatrix}.$$

The eigenvalues of A are

$$\begin{aligned} & -0.0640196 + 0.732597i, & -0.0640196 - 0.732597i, \\ & -0.390646 + 0.477565i, & -0.390646 - 0.477565i, \\ & -0.756193 + 0.125533i, & -0.756193 - 0.125533i. \end{aligned}$$




FIG. 6.7. *Example 4*, $n = 6, k = 2$, A normal real, location of $\theta_1 = \bar{\theta}_2$

Figure 6.8 is an example with two pairs, one real eigenvalue on the boundary of the field of values and one real eigenvalue inside (Example 5). We have two spurious curves. The matrix A is

$$A = \begin{pmatrix} -0.500411 & 0.25411 & 0.499092 & -0.15696 & 1.26376 & -0.690147 \\ -0.850536 & 0.662412 & 0.12518 & 0.666057 & -0.873974 & -0.503358 \\ -0.095158 & 0.54861 & -0.0510311 & -0.42028 & -0.209823 & 0.122187 \\ 0.307198 & 0.827682 & -0.341422 & -0.437352 & 0.0411078 & -0.835649 \\ -1.00153 & 0.456062 & -0.0256999 & -0.551469 & 0.191305 & 1.01331 \\ 0.762756 & 0.970216 & 0.404506 & 0.804347 & 0.368779 & 0.630639 \end{pmatrix}.$$

The eigenvalues of A are

$$\begin{aligned} & -0.432565 + 1.66558i, & -0.432565 - 1.66558i, \\ & 1.19092 + 1.18916i, & 1.19092 - 1.18916i, \\ & -1.20852, & 0.187377. \end{aligned}$$

To visualize the feasible region for θ_1 it could be good to get rid of the “spurious” curves. This can be done approximately in the following way. We can compute points on the curves by solving an equation in x for a given value of y (or vice-versa) for each equation defining the boundary. When we get a point on a curve, we can check points surrounding it in the complex plane. If there is at least one of those points which is not feasible then our given point is on the boundary and the piece of the curve on which it is located is a part of the boundary. This is not always easy because of rounding errors and because we could have some curves which are almost tangent to each other. Of course this process is not foolproof since the result depends on the choice of the surrounding points and also on some thresholds. But in many cases it works fine. Figure 6.9 shows what we get for the previous example. The blue stars




FIG. 6.8. Example 5, $n = 6, k = 2$, A normal real, location of $\theta_1 = \bar{\theta}_2$

are the points computed on the boundary (using the Matlab function `fzero`). Note that we get rid of the two spurious curves since we keep only the curves on which there is at least one boundary point.




FIG. 6.9. Example 5, $n = 6, k = 2$, boundary of the feasible region

There is another way to visualize the boundary of the feasible region in Matlab. The `contour` function that we use is evaluating the function on a grid and then finding the curve of level 0 by interpolation. Therefore, we can set up a routine that given x and y computes a solution of the underdetermined system for the point $(x + iy, x - iy)$ using the SVD. If the point is not feasible then we return a very small negative value. However, this process is very expensive since the evaluation of the function cannot be vectorized. An example is given below in Figure 6.11 for the next Example 6. Of course we do not have spurious curves and not even the parts of the curves that are not relevant. But we have some wiggles in the curve because we set the values for non-feasible points to a small negative value introducing discontinuities in the function values.

Figure 6.10 displays an example with two pairs (one inside the FOV) and two real eigenvalues (Example 6). The feasible region has an interesting shape. Figure 6.11 shows the boundary for Example 6 computed using the SVD. The matrix is

$$A = \begin{pmatrix} 0.0433091 & 1.59759 & -0.318964 & -0.787924 & -1.5765 & 0.538701 \\ 0.222478 & -0.276959 & 0.775185 & 1.54146 & 1.8561 & 0.818277 \\ 0.348846 & -0.0614769 & 1.02246 & -0.677541 & -0.498161 & 0.193331 \\ 1.05979 & -1.7532 & -0.176368 & 0.214925 & -0.563343 & -0.580403 \\ 2.01859 & -0.900034 & 0.21777 & -1.05788 & -0.388673 & -1.0512 \\ 0.825456 & -0.837442 & 0.298154 & -0.554189 & 0.812614 & 0.77613 \end{pmatrix}.$$

The eigenvalues of A are

$$\begin{aligned} & -1.2413 + 3.27037i, & -1.2413 - 3.27037i, \\ & 0.566382 + 0.768588i, & 0.566382 - 0.768588i, \\ & 0.917215 & 1.82382. \end{aligned}$$




FIG. 6.10. Example 6, $n = 6, k = 2$, A normal real, location of $\theta_1 = \bar{\theta}_2$




FIG. 6.11. Example 6, $n = 6, k = 2$, A normal real, boundary obtained with the SVD

Figure 6.12 displays an example with $n = 8$ (Example 7). We can see that (unfortunately) we have many spurious curves that are useless for the boundary. On

the right part of Figure 6.12 we got rid of some of these curves but not all of them. The matrix of Example 7 is

$$A = \begin{pmatrix} 0.541379 & 0.36045 & 0.724658 & -0.835226 & -0.882172 & 0.0513467 & -0.231744 & -0.316297 \\ -0.454221 & 0.575524 & -0.100099 & -0.312607 & -0.365987 & -0.122991 & 0.143776 & 0.447837 \\ 0.210676 & -0.0931479 & 0.852157 & 0.39926 & -0.119268 & -0.722606 & 0.199469 & 0.255216 \\ -0.921745 & -0.357353 & 0.0571532 & -0.569208 & -1.24529 & 1.17068 & 0.120452 & -0.304355 \\ -0.719429 & -0.137593 & 0.470774 & -1.33238 & -0.162772 & 1.02581 & -0.277858 & 0.154487 \\ 0.451727 & 0.489061 & -0.0903518 & 0.835521 & 1.06541 & 0.646274 & -0.158683 & 0.856737 \\ 0.00891101 & 0.0305841 & -0.23076 & -0.0649839 & 0.0463489 & 0.236475 & 0.810799 & -0.356549 \\ 0.682085 & -0.398763 & 0.179775 & -0.759383 & -0.268957 & 0.158633 & -0.112359 & 1.03084 \end{pmatrix}.$$

The eigenvalues of A are

$$\begin{aligned} &1.68448 + 0.780709i, & 1.68448 - 0.780709i, \\ &0.418673 + 0.888289i, & 0.418673 - 0.888289i, \\ &0.882938 + 0.19178i, & 0.882938 - 0.19178i, \\ &-2.9958, & 0.748615. \end{aligned}$$




FIG. 6.12. Example 7, $n = 8, k = 2$, A normal real, location of $\theta_1 = \bar{\theta}_2$

We remark that using the same technique as before we can compute the boundary of the feasible region for θ_2 when θ_1 is prescribed for $k = 2$ and complex normal matrices. Here we have to consider basic solutions for the real matrix which is $5 \times n$. Hence we compute the solutions for all 5×5 matrices extracted from the system for ω . In this case we compute the solutions numerically and not symbolically for a given (x, y) . Then we check that the curves are really parts of the boundary using the same perturbation technique as before. We consider the problem of Figure 1 (b) in Bujanović [3]. The eigenvalues of A are $-5, -3 + 2i, -3 - 2i, 4 + i, 4 - i, 6$. We fix $\theta_1^{(2)} = \theta_1 = -4$. The boundary of the feasible region for θ_2 for this particular value of θ_1 is displayed in Figure 6.13.

One can compare with [3] and see that we indeed find the boundary of the region for θ_2 . However, such regions do not give a good idea of the locations of the Ritz values because we would have to move θ_1 all over the field of values to see where the Ritz values can be located. Figure 6.14 displays the locations of the Ritz values for $k = 2$ to 5 when running Arnoldi with a complex diagonal matrix with the given eigenvalues and random real starting vectors. We see that we have Ritz values almost everywhere. Things are strikingly different if we construct a real normal matrix with the given eigenvalues (which are real or occur in complex conjugate pairs) and run Arnoldi with real starting vectors. The Ritz values are shown in Figure 6.15. We see that they are constrained in two regions of the complex plane and on the real axis. Of course things would have been different if we would have used complex starting vectors. The Ritz values would have looked more like in Figure 6.14. There is much more structure in the feasible region if everything is real.




FIG. 6.13. Example in Figure 1(b) of [3], $n = 6, k = 2, \theta_1^{(2)} = -4$, boundary of the feasible region for $\theta_2^{(2)}$




FIG. 6.14. $n = 6$, all $k = 2 : 5$, A complex diagonal, location of the Ritz values, Arnoldi with random real vectors v

7. Open problems and conjectures for $k > 2$ and real normal matrices.

In this section we describe some numerical experiments with $k > 2$ for real normal matrices. We also state some open problems and we formulate some conjectures. We are interested in $k = 3$ to $k = n - 1$. We would like to know where are the Ritz values located when using real starting vectors. Clearly we cannot do the same as for $k = 2$ because, for instance, for $k = 3$ we have either three real Ritz values or a pair of complex conjugate Ritz values and a real one. Of course, we can fix the location of the real Ritz values and look for the region where the pairs of the complex conjugate Ritz values may be located but this is not that informative since it is not practical to explore all the possible locations of the real Ritz values.

Let us do some numerical experiments with random starting vectors and first consider Example 6 of the previous subsection with $n = 6$. For each value of $k = 2 : n - 1$ we generate 700 random initial vectors of unit norm and we run the Arnoldi




FIG. 6.15. $n = 6$, all $k = 2 : 5$, A normal real, location of the Ritz values, Arnoldi with random real vectors v

algorithms computing the Ritz values at iteration k . In Figure 7.1 we plot the pairs in blue and red and the real eigenvalues in green for all the values of k and we superimpose the boundary curves computed for $k = 2$. We see that all the Ritz values belong to the feasible region that was computed for $k = 2$. We conjecture that this is true for any real normal matrix and a real starting vector. But there is more than that.

Figure 7.2 displays the Ritz values at iteration 4. We see that some of the Ritz values are contained in a region for which one part of the boundary is one piece of a curve that was considered as “spurious” for $k = 2$. Figure 7.3 shows the Ritz values at iteration 5 (that is, the next to last one); there is an accumulation of some Ritz values on this spurious curve as well as close to the other pair of complex conjugate eigenvalues. It seems that some of the spurious curves look like “attractors” for the Ritz values, at least for random real starting vectors. It would be interesting to explain this phenomenon.




FIG. 7.1. Example 6, $n = 6$, all $k = 2 : 5$, A normal real, location of the Ritz values, Arnoldi with random real vectors v




FIG. 7.2. *Example 6, $n = 6$, $k = 4$, A normal real, location of the Ritz values, Arnoldi with random real vectors v*




FIG. 7.3. *Example 6, $n = 6$, $k = 5$, A normal real, location of the Ritz values, Arnoldi with random real vectors v*

Figures 7.4 to 7.8 give the results for Example 7 with $n = 8$. Here again we see that the Ritz values are inside the boundary for $k = 2$ and, at some iterations, Ritz values are located preferably on or close to some of the spurious curves.

Another open question is to know if there exist real normal matrices for which the feasible region for $k = 2$ completely fill the field of values for real starting vectors. In this paper we concentrated on pairs of complex conjugate Ritz values but an interesting problem is to locate the real Ritz values in the intersection of the field of values with the real axis.

Numerical experiments not reported here seem to show that the properties described above for the Arnoldi Ritz values are not restricted to the Arnoldi algorithm. For a real normal matrix, if one constructs a real orthogonal matrix V and define $H = V^T A V$, the Ritz values, being defined as the eigenvalues of H_k the principal submatrix of order k of H , are also constrained in some regions inside the field of




FIG. 7.4. Example 7, $n = 8$, all $k = 2 : 7$, A normal real, location of the Ritz values, Arnoldi with random real vectors v




FIG. 7.5. Example 7, $n = 8$, $k = 4$, A normal real, location of the Ritz values, Arnoldi with random real vectors v

values of A . This deserves further studies.

8. Conclusion. In this paper we gave a necessary and sufficient condition for a set of complex values $\theta_1, \dots, \theta_k$ to be the Arnoldi Ritz values at iteration k for a general diagonalizable matrix A . This generalizes previously known conditions. This condition simplifies for normal matrices and particularly for real normal matrices and real starting vectors. We particularly studied the case $k = 2$ for which we characterized the boundary of the region in the complex plane contained in $W(A)$ where pairs of complex conjugate Ritz values are located. Several examples of computation of the boundary of the feasible region were given. Finally, after describing some numerical experiments with random real starting vectors, we formulated some conjectures and open problems for $k > 2$ for real normal matrices.

Acknowledgments. The author thanks J. Duintjer Tebbens for some interesting comments.




FIG. 7.6. Example 7, $n = 8$, $k = 5$, A normal real, location of the Ritz values, Arnoldi with random real vectors v




FIG. 7.7. Example 7, $n = 8$, $k = 6$, A normal real, location of the Ritz values, Arnoldi with random real vectors v

REFERENCES

- [1] M. BELLALIJ, Y. SAAD AND H. SADOK, *On the convergence of the Arnoldi process for eigenvalue problems*, Matrix Analysis and Applications, (2007), pp. 5.
- [2] M. BELLALIJ, Y. SAAD AND H. SADOK, *Further analysis of the Arnoldi process for eigenvalue problems*, SIAM J. Numer. Anal., v 48 n 2 (2010), pp. 393–407.
- [3] Z. BUJANOVIĆ, *On the permissible arrangements of Ritz values for normal matrices in the complex plane*, Linear Algebra Appl., v 438 (2013), pp. 4606–4624.
- [4] R. CARDEN, *Ritz values and Arnoldi convergence for non-Hermitian matrices*, Ph.D. thesis, Rice University (2011).
- [5] R. CARDEN, *A simple algorithm for the inverse field of values problem*, Inverse Problems, v 25 (2009), pp. 1–9.
- [6] R. CARDEN AND M. EMBREE, *Ritz value localization for non-Hermitian matrices*, SIAM J. Matrix Anal. Appl., v 33 (2012), pp. 1320–1338.
- [7] R. CARDEN AND D.J. HANSEN, *Ritz values of normal matrices and Ceva's theorem*, Linear Algebra Appl., v 438 (2013), pp. 4114–4129.
- [8] C. CHORIANOPOULOS, P. PSARRAKOS AND F. UHLIG, *A method for the inverse numerical range*




FIG. 7.8. Example 7, $n = 8$, $k = 7$, A normal real, location of the Ritz values, Arnoldi with random real vectors v

- problem, *Elec. J. Linear Alg.*, v 20 (2010), pp. 198–206.
- [9] J. DUINTJER TEBBENS AND G. MEURANT, *Any Ritz value behavior is possible for Arnoldi and for GMRES*, *SIAM J. Matrix Anal. Appl.*, v 33 n 3 (2012), pp. 958–978.
- [10] J. DUINTJER TEBBENS AND G. MEURANT, *On the convergence of QOR and QMR Krylov methods for solving linear systems*, in preparation (2013).
- [11] I.C.F. IPSEN, *Expressions and bounds for the GMRES residual*, *BIT*, v 40 n 3 (2000), pp. 524–535.
- [12] A. KOVAČEC AND B. RIBEIRO, *Convex hull calculations: A Matlab implementation and correctness proofs for the lrs-algorithm*, technical report, Coimbra University, Portugal (2003).
- [13] R.B. LEHOUCQ, D.C. SORENSEN, AND CHAO-CHIH YANG, *Arpack User's Guide: solution of large-scale eigenvalue problems with implicitly restarted Arnoldi methods*, Vol. 6 Siam, (1998).
- [14] S.M. MALAMUD, *Inverse spectral problem for normal matrices and the Gauss-Lucas theorem*, *Transactions of the AMS*, v 357 n 10 (2004), pp. 4043–4064.
- [15] G. MEURANT, *GMRES and the Arioli, Pták, and Strakoš parametrization*, *BIT Numerical Mathematics*, v 52 n 3 (2012), pp. 687–702.
- [16] G. MEURANT, *The computation of isotropic vectors*, *Numerical Algorithms*, v 60 n 2 (2012), pp. 193–204.
- [17] C.C. PAIGE, *The computation of eigenvalues and eigenvectors of very large sparse matrices*, Ph.D. thesis, University of London, (1971).
- [18] C.C. PAIGE, *Computational variants of the Lanczos method for the eigenproblem*, *J. Inst. Maths Applics.*, v 10, (1972), pp 373–381.
- [19] C.C. PAIGE, *Accuracy and effectiveness of the Lanczos algorithm for the symmetric eigenproblem*, *Linear Algebra Appl.*, v 34, (1980), pp 235–258.
- [20] B.N. PARLETT, *Normal Hessenberg and moment matrices*, *Linear Algebra Appl.*, v 6 (1973), pp. 37–43.
- [21] B.N. PARLETT, *The symmetric eigenvalue problem*, Prentice Hall, (1980). Reprinted by SIAM in the series *Classics in Applied Mathematics* (1998).
- [22] H. SADOK, *Analysis of the convergence of the minimal and the orthogonal residual methods*, *Numerical Algorithms*, v 40 (2005), pp. 201–216.